

University of Groningen

Towards adipose tissue-derived stromal cells-based therapy for diabetic retinopathy

Hajmoussa, Ghazaleh

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Hajmoussa, G. (2018). *Towards adipose tissue-derived stromal cells-based therapy for diabetic retinopathy*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Towards adipose tissue-derived stromal cells-based therapy for diabetic retinopathy

Propositions:

1. Adipose tissue derived stromal cells (ASC) support newly formed vascular networks by endothelial cells (EC) in vitro and in vivo through maintenance of the vascular architecture (this thesis).
2. Administered ASC to the murine model of diabetic retinopathy augment and stabilize retinal angiogenesis and co-localize with capillaries at a pericyte-specific position (this thesis).
3. The ROS-induced mitochondrial dysfunction, hyperglycemia-induced apoptosis, and bioenergetics changes only partially influence the pericytic ability of ASC (this thesis).
4. Despite of the diminished level of glycolysis, ASC under HG showed the same basal respiration as under NG which proves that ASC resist the HG condition by changing the metabolic limit deprived of altering the proliferation rate (this thesis).
5. Conditioned medium of ASC alleviates high glucose-induced oxidative stress and its subsequent upregulated downstream targets in an NF- κ B dependent fashion in bovine retinal endothelial cells (this thesis).
6. The manner of high glucose preconditioning of adipose tissue-derived stromal cells dictates their immuno-regulatory properties (this thesis).
7. The antioxidant capacity of ASC-Cme downmodulates upregulated pro-inflammatory genes in high glucose-challenged BREC (this thesis).
8. The new pharmacologic compound, (6-hydroxyl-2,5,7,8-tetramethylchroman-2-yl)(4-(2-hydroxyethyl)piperazin-1-yl) methanone (SUL-109) shields ASC during cell preservation from hypothermic cell death without influencing their multi-potency capacity and proliferation (this thesis).